

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use SINCALIDE FOR INJECTION safely and effectively. See full prescribing information for SINCALIDE FOR INJECTION

SINCALIDE for injection, for intravenous use  
Initial U.S. Approval: 1976

### RECENT MAJOR CHANGES

Contraindications (4) 10/2023

Warnings and Precautions (5.1)

Anaphylaxis, Anaphylactic Shock and Other Serious Hypersensitivity Reactions (5.1) 10/2023

### INDICATIONS AND USAGE

Sinacalide for Injection is a cholecystokinin (CCK) analog indicated in adults to:

- stimulate gallbladder contraction, as may be assessed by various methods of diagnostic imaging, or to obtain by duodenal aspiration a sample of concentrated bile for analysis of cholesterol, bile salts, phospholipids, and crystals. (1)
- stimulate pancreatic secretion in combination with secretin prior to obtaining a duodenal aspirate for analysis of enzyme activity, composition, and cytology. (1)
- accelerate the transit of a barium meal through the small bowel, thereby decreasing the time and extent of radiation associated with fluoroscopy and x-ray examination of the intestinal tract. (1)

### DOSAGE AND ADMINISTRATION

Recommended Adult Dosage and Administration by Indication:

#### To Stimulate Contraction of the Gallbladder

- 0.02 mcg/kg as a single dose over 30 to 60 seconds via intravenous injection. If satisfactory contraction does not occur in 15 minutes, administer a dose of 0.04 mcg/kg over 30 to 60 seconds. (2.1)

- Alternatively consider an intravenous infusion to reduce gastrointestinal adverse reactions:* 0.12 mcg/kg diluted in 100 mL of 0.9% Sodium Chloride Injection USP and infused over 50 minutes at a rate of 2 mL per minute. (2.1, 2.2, 5.3)

#### To Stimulate Pancreatic Secretion in Combination with Secretin

- 30 minutes after initiation of secretin for injection, administer 0.02 mcg/kg diluted in 30 mL of 0.9% Sodium Chloride Injection USP and infused over 30 minutes at a rate of 1 mL per minute. (2.1, 2.2)

#### To Accelerate Transit of a Barium Meal Through the Small Intestine

- After the barium meal is beyond the proximal jejunum, administer 0.04 mcg/kg over 30 to 60 seconds via intravenous injection. (2.1)

- If satisfactory transit of the barium meal has not occurred in 30 minutes, administer a second dose of 0.04 mcg/kg over 30 to 60 seconds. (2.1)
- Alternatively consider an intravenous infusion to reduce gastrointestinal adverse reactions:* 0.12 mcg/kg diluted in 100 mL 0.9% Sodium Chloride Injection USP and infused over 30 minutes. (2.1, 2.2, 5.3)

### DOSAGE FORMS AND STRENGTHS

For injection: 5 mcg of sinacalide as a lyophilized powder in a single-dose vial for reconstitution (3)

### CONTRAINDICATIONS

- History of hypersensitivity to sulfites or sinacalide. (4, 5.1)
- Intestinal obstruction. (4)

### WARNINGS AND PRECAUTIONS

- Anaphylaxis, Anaphylactic Shock and Other Serious Hypersensitivity Reactions:** Contains sodium metabisulfite. Serious reactions may occur during or soon after administration. If symptoms occur, discontinue the drug. (4, 5.1)
- Evacuation of Gallstones:** Stimulation of gallbladder contraction in patients with small gallbladder stones could lead to the evacuation of the stones from the gallbladder, resulting in their lodging in the cystic duct or in the common bile duct. (5.2)
- Gastrointestinal Adverse Reactions with Intravenous Injection:** Administration as an intravenous injection may cause transient nausea, vomiting, abdominal pain or cramping, dizziness or flushing. To reduce the risk of adverse reactions when used to stimulate contraction of the gallbladder or accelerate transit of a barium meal through the small intestine, administer as an intravenous infusion over 50 or 30 minutes, respectively. (2.1, 5.3)
- Preterm Labor or Spontaneous Abortion:** Advise pregnant women of the potential risk for preterm labor and spontaneous abortion. (5.4, 8.1)

### ADVERSE REACTIONS

Most common adverse reactions (≥20%) are: abdominal discomfort or pain, and nausea. (6)  
**To report SUSPECTED ADVERSE REACTIONS, contact Fresenius Kabi USA, LLC at 1-800-551-7176 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.**

### DRUG INTERACTIONS

**Drugs that Affect Gallbladder Motility or Contractile Response:** May interfere with response to sinacalide. Consider discontinuing these drugs prior to administration of Sinacalide for Injection, when used to stimulate contraction of the gallbladder. (7.1)

**See 17 for PATIENT COUNSELING INFORMATION.**

Revised: 10/2023

## FULL PRESCRIBING INFORMATION: CONTENTS\*

### 1 INDICATIONS AND USAGE

#### 2 DOSAGE AND ADMINISTRATION

- Recommended Dosage and Administration Instructions
- Preparation Instructions

#### 3 DOSAGE FORMS AND STRENGTHS

#### 4 CONTRAINDICATIONS

#### 5 WARNINGS AND PRECAUTIONS

- Anaphylaxis, Anaphylactic Shock and Other Serious Hypersensitivity Reactions
- Evacuation of Gallstones
- Gastrointestinal Adverse Reactions with Intravenous Injection
- Preterm Labor or Spontaneous Abortion

#### 6 ADVERSE REACTIONS

## FULL PRESCRIBING INFORMATION

### 1 INDICATIONS AND USAGE

Sinacalide for Injection is indicated in adults to:

- to stimulate gallbladder contraction, as may be assessed by various methods of diagnostic imaging, or to obtain by duodenal aspiration a sample of concentrated bile for analysis of cholesterol, bile salts, phospholipids, and crystals;
- to stimulate pancreatic secretion in combination with secretin prior to obtaining a duodenal aspirate for analysis of enzyme activity, composition, and cytology;

### 7 DRUG INTERACTIONS

- Drugs that Affect Gallbladder Motility or Contractile Response

### 8 USE IN SPECIFIC POPULATIONS

- Pregnancy
- Lactation
- Pediatric Use
- Geriatric Use

### 10 OVERDOSAGE

### 11 DESCRIPTION

### 12 CLINICAL PHARMACOLOGY

- Mechanism of Action
- Pharmacodynamics

### 13 NONCLINICAL TOXICOLOGY

- Carcinogenesis, Mutagenesis, Impairment of Fertility

### 16 HOW SUPPLIED/STORAGE AND HANDLING

### 17 PATIENT COUNSELING INFORMATION

\* Sections or subsections omitted from the full prescribing information are not listed.

- to accelerate the transit of a barium meal through the small bowel, thereby decreasing the time and extent of radiation associated with fluoroscopy and x-ray examination of the intestinal tract.

### 2 DOSAGE AND ADMINISTRATION

#### 2.1 Recommended Dosage and Administration Instructions

The recommended dosage and administration of Sinacalide for Injection by indication is shown in Table 1. For preparation instructions see *Dosage and Administration* (2.2).

## Table 1: Recommended Adult Dosage and Administration of Sinacalide for Injection by Treatment Indication

Indication	Recommended Adult Dosage and Administration of Sinacalide for Injection
To stimulate contraction of the gallbladder	Sinacalide for Injection 0.02 mcg/kg as a single dose over 30 to 60 seconds via intravenous injection. If satisfactory contraction does not occur in 15 minutes, administer a dose of 0.04 mcg/kg over 30 to 60 seconds. <i>Alternatively, Consider an Intravenous Infusion to Reduce Gastrointestinal Adverse Reactions. [See Warnings and Precautions (5.3)]:</i> 0.12 mcg/kg diluted in 100 mL of 0.9% Sodium Chloride Injection USP and infused over 50 minutes at a rate of 2 mL per minute.
To stimulate pancreatic secretion in combination with secretin for injection	Secretin for Injection: 0.25 units/kg as intravenous infusion over 60 minutes Sinacalide for Injection: 30 minutes after initiation of secretin infusion, administer Sinacalide for Injection 0.02 mcg/kg diluted in 30 mL of 0.9% Sodium Chloride Injection USP and infused over 30 minutes at a rate of 1 mL per minute.
To accelerate the transit of a barium meal through the small intestine	After the barium meal is beyond the proximal jejunum, administer Sinacalide for Injection 0.04 mcg/kg over 30 to 60 seconds via intravenous injection. If satisfactory transit of the barium meal has not occurred in 30 minutes, administer a second dose of 0.04 mcg/kg over 30 to 60 seconds. <i>Alternatively, Consider an Intravenous Infusion to Reduce Gastrointestinal Adverse Reactions. [See Warnings and Precautions (5.3)]:</i> 0.12 mcg/kg diluted in 100 mL 0.9% Sodium Chloride Injection USP and infused over 30 minutes.

### 2.2 Preparation Instructions

#### For Intravenous Injection

- Reconstitute Sinacalide for Injection aseptically by adding 5 mL of Sterile Water for Injection USP to the vial.
- Inspect the reconstituted solution visually for particulate matter and discoloration after reconstitution and prior to administration.
- Withdraw the prescribed dose of the reconstituted solution from the vial and administer as an intravenous injection over 30 to 60 seconds, as shown in Table 1. Discard the unused portion.
- Store the reconstituted solution at room temperature. *Discard after 8 hours.*
- For single use only; discard unused portion.

#### For Intravenous Infusion

- Reconstitute Sinacalide for Injection aseptically by adding 5 mL of Sterile Water for Injection USP to the vial.
- After reconstitution, withdraw the prescribed dose of the solution from the vial. Discard unused portion.
- Dilute the reconstituted solution in 30 mL or 100 mL of 0.9% Sodium Chloride Injection USP, depending on the indication, as described in Table 1.
- Inspect the Sinacalide for Injection solutions visually for particulate matter and discoloration after reconstitution, dilution and prior to administration.
- Store the diluted solution at room temperature. *Discard after 1 hour.*

### 3 DOSAGE FORMS AND STRENGTHS

For injection: 5 mcg of sinacalide as a lyophilized white powder for reconstitution in a single-dose vial.

### 4 CONTRAINDICATIONS

Sinacalide for Injection is contraindicated in patients with:

- a history of hypersensitivity to sulfites or sinacalide. Serious hypersensitivity reactions have included anaphylaxis and anaphylactic shock [see *Warnings and Precautions* (5.1), *Adverse Reactions* (6)].
- intestinal obstruction.

### 5 WARNINGS AND PRECAUTIONS

#### 5.1 Anaphylaxis, Anaphylactic Shock and Other Serious Hypersensitivity Reactions

Contains sodium metabisulfite [see *Description* (11)], a sulfite that may cause allergic-type reactions including anaphylactic symptoms and life-threatening or less severe asthmatic episodes in certain susceptible people. The overall prevalence of sulfite sensitivity in the general population is unknown and probably low. Sulfite sensitivity is seen more frequently in asthmatic than in non-asthmatic people.

In postmarketing experience, anaphylaxis, anaphylactic shock and other serious hypersensitivity reactions have been reported during and within one hour following administration of Sinacalide for Injection [see *Adverse Reactions* (6)].

Sinacalide for Injection is contraindicated in patients with a history of hypersensitivity to sulfites. Due to the potential for anaphylaxis, appropriate medical support should be readily available when Sinacalide for Injection is administered. If anaphylaxis

or other hypersensitivity reactions occur, immediately discontinue the infusion and initiate appropriate medical treatment. Observe patients closely during and after the infusion. Do not reinitiate Sinacalide for Injection in patients who have experienced symptoms of hypersensitivity [see *Contraindications* (4)].

### 5.2 Evacuation of Gallstones

Stimulation of gallbladder contraction in patients with small gallbladder stones could lead to the evacuation of the stones from the gallbladder, resulting in their lodging in the cystic duct or in the common bile duct.

### 5.3 Gastrointestinal Adverse Reactions with Intravenous Injection

Administration of Sinacalide for Injection as an intravenous injection may cause adverse reactions such as nausea, vomiting, abdominal pain or cramping, dizziness, and flushing [see *Adverse Reactions* (6)]. These reactions are generally transient. To reduce the risk of adverse reactions with intravenous injection when used to stimulate contraction of the gallbladder or accelerate transit of a barium meal through the small intestine, administer Sinacalide for Injection as an intravenous infusion over 50 or 30 minutes, respectively [see *Dosage and Administration* (2.1)].

### 5.4 Preterm Labor or Spontaneous Abortion

Because of Sinacalide for Injection's effect on smooth muscle, pregnant patients should be advised that spontaneous abortion or premature induction of labor may occur [see *Use in Specific Populations* (8.1)].

## 6 ADVERSE REACTIONS

The following clinically significant adverse reactions are described elsewhere in labeling:

- Anaphylaxis, anaphylactic shock, and other serious hypersensitivity reactions [see *Warnings and Precautions* (5.1)]
- Evacuation of gallstones [see *Warnings and Precautions* (5.2)]
- Adverse reactions with intravenous injection [see *Warnings and Precautions* (5.3)]
- Preterm labor or spontaneous abortion [see *Warnings and Precautions* (5.4)]

The following adverse reactions associated with the use of Sinacalide for Injection were identified in clinical trials or postmarketing reports. Because these reactions were reported voluntarily from a population of uncertain size, it is not always possible to estimate their frequency, reliably, or to establish a causal relationship to drug exposure.

The most frequent adverse reactions (20% or greater) are gastrointestinal: abdominal discomfort or pain, and nausea; these may not necessarily indicate an abnormality of the biliary tract unless there is other clinical or radiologic evidence of disease.

Less common adverse reactions include:

**Hypersensitivity reactions:** anaphylaxis and anaphylactic shock, hypotension, throat tightness, bradycardia, shortness of breath, nausea, abdominal cramping, diaphoresis, hives, rash, itching; and numbness of face, lips and eyes [see *Contraindications* (4), (5.1)].

**Neurological reactions:** seizures, headache.

**Vasovagal reactions:** dizziness, loss of consciousness, nausea, diaphoresis, syncope and hypotension (generally self-limiting).

**Other:** nausea, vomiting, flushing, hypertension, urge to defecate, diarrhea, sneezing.

### 7 DRUG INTERACTIONS

#### 7.1 Drugs that Affect Gallbladder Motility or Contractile Response

Drugs that may stimulate or inhibit gallbladder motility or contractile response may interfere with the response to sinacalide. Consider discontinuing these drugs prior to administration of Sinacalide for Injection, when used to stimulate contraction of the gallbladder.

### 8 USE IN SPECIFIC POPULATIONS

#### 8.1 Pregnancy

##### Risk Summary

Based on limited human data and mechanism of action, sinacalide may cause preterm labor or spontaneous abortion [see *Warnings and Precautions* (5.4)]. Available data with Sinacalide for Injection are insufficient to establish a drug-associated risk of major birth defects, miscarriage or adverse maternal or fetal outcomes. In animal embryo-fetal development studies in which sinacalide was administered to hamsters and rats during the period of organogenesis, no effects were seen at doses comparable to the maximum recommended clinical dose on a mg/kg basis. However, in a prenatal development study in which rats were administered sinacalide during organogenesis through parturition, decreased weight gain and developmental delays were observed at a dose 122 times higher than the maximum recommended human dose based on body surface area.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

#### Data

##### Animal Data

There were no effects on embryo-fetal development in hamsters when sinacalide was administered subcutaneously at 250 or 750 ng/kg during organogenesis (Gestation Days 7 to 13) at doses up to 0.8 times the maximum recommended dose of 120 ng/kg on a body surface area basis. No effects on embryo-fetal development were observed in Sprague-Dawley rats at subcutaneous doses of 250, 450, or 750 ng/kg from Gestation Days 6 to 16, representing 1.0 time the maximum recommended human dose on a body surface area basis. In a separate study at a higher dose of 90 mcg/kg administered subcutaneously to CFY rats from Gestation Day 10 through parturition (representing 122 times the maximum recommended human dose on a body surface area basis), offspring showed decreased growth, behavioral changes, and developmental delays.

### 8.2 Lactation

#### Risk Summary

There are no data regarding the presence of sinacalide in human or animal milk, the effects on the breastfed infant, or the effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Sinacalide for Injection and any potential adverse effect on the breastfed infant from Sinacalide for Injection or from the underlying condition.

### 8.4 Pediatric Use

The safety and effectiveness in pediatric patients have not been established.

### 8.5 Geriatric Use

Clinical studies of Sinacalide for Injection did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects.

## 10 OVERDOSAGE

In the event of an overdose, symptoms related to vagal stimulation, such as gastrointestinal symptoms (abdominal cramps, nausea, vomiting and diarrhea), hypotension with dizziness or fainting may occur. Overdosage symptoms should be treated symptomatically and should be of short duration.


A single bolus intravenous injection of 0.05 mcg/kg (approximately 2 to 3 times the human dose of 0.02 mcg/kg), sinacalide caused hypotension and bradycardia in dogs. In addition, higher doses injected intravenously once or repeatedly in dogs caused syncope and ECG changes (approximately 5 times the human dose of 0.02 mcg/kg). These effects were attributed to sinacalide-induced vagal stimulation in that all were prevented by pretreatment with atropine or bilateral vagotomy.

## 11 DESCRIPTION

Sinacalide for Injection is a cholecystopancreatic-gastrointestinal hormone for parenteral administration. The agent is a synthetically-prepared C-terminal octapeptide of cholecystokinin.

Each single-dose vial of sinacalide provides a sterile nonpyrogenic lyophilized white powder consisting of 5 mcg sinacalide with 30 mg arginine hydrochloride, 15 mg lysine hydrochloride, 170 mg mannitol, 4 mg methionine, 2 mg pentetic acid, 0.005 mcg polysorbate 20, 9 mg potassium phosphate dibasic, and 0.04 mg sodium metabisulfite.

The pH is adjusted to 6.0 to 8.0 with hydrochloric acid and/or sodium hydroxide prior to lyophilization. Sinacalide is designated chemically as L- $\alpha$ -aspartyl-O-sulfo-L-tyrosyl-L-methionylglycyl-L-tryptophyl-L-methionyl-L- $\alpha$ -aspartyl-L-phenylalaninamide. Graphic formula:



## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

When injected intravenously, sinacalide stimulates gallbladder contraction and reduction in size. The evacuation of bile that results is similar to that which occurs physiologically in response to endogenous cholecystokinin. Sinacalide also stimulates pancreatic secretion and intestinal motility causing pyloric contraction and slows gastric emptying.



4 51763A /Revised: October 2023

# SINCALIDE for Injection

Rx only

Concurrent administration of sincalide with secretin increases both the volume of pancreatic secretion and the out-put of bicarbonate and enzymes. This combined effect of secretin and sincalide permits the assessment of specific pancreatic function through measurement and analysis of the duodenal aspirate.

## 12.2 Pharmacodynamics

Following an intravenous (bolus) injection of 0.02 mcg/kg of sincalide, maximal contraction of the gallbladder occurred in 5 to 15 minutes. Sincalide reduced gallbladder radiographic size by at least 40%, which is generally considered satisfactory contraction.

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term studies in animals have not been performed to evaluate carcinogenic or mutagenic potential, or possible impairment of fertility in males or females.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

Sincalide for Injection is supplied as 5 mcg of sincalide as a lyophilized white powder for reconstitution in a single-dose vial; in packages of 10 vials as follows:

Product Code	Unit of Sale	Strength	Each
579105	NDC 63323-579-05 Unit of 10	5 mcg per vial	NDC 63323-579-01 Single-Dose Vial

Store at 25° C (77° F); excursions permitted to 15-30° C (59-86° F) [See USP Controlled Room Temperature].

## 17 PATIENT COUNSELING INFORMATION

### Anaphylaxis, Anaphylactic Shock and Other Serious Hypersensitivity Reactions

Inform patients that serious hypersensitivity reactions, including anaphylaxis and anaphylactic shock have been reported during or following administration of Sincalide for Injection. Advise patients to report immediately to a healthcare provider if they experience symptoms of a hypersensitivity reaction [see Warnings and Precautions (5.1)].

### Gastrointestinal Adverse Reactions

Advise patients that Sincalide for Injection may cause transient gastrointestinal symptoms [see Warnings and Precautions (5.3)].

### Pregnancy

Advise pregnant women of the potential risk for preterm labor and spontaneous abortion [see Warnings and Precautions (5.4), Use in Specific Populations (8.1)].

## Rx only

Manufactured by:



Lake Zurich, IL 60047  
www.fresenius-kabi.com/us